

## Summer School in Glaciology, McCarthy, 2022

**Exercises: Glacier meteorology / Energy balance****1.) ENERGY BALANCE – MELT CALCULATION**

Assume a summer melt season and the glacier **surface is at melting point** ( $0^{\circ}\text{C}$ ), but the glacier is polythermal with a **cold surface layer**, i.e. below the surface there is a layer where the glacier's temperature is below freezing and therefore energy is consumed not only for melting but also to warm up that layer. The following variables (daily means) are measured at a weather station on the glacier. Heat fluxes providing energy to the surface are positive; heat fluxes away from the surface are negative (consuming energy). Energy sources are positive, energy sinks are negative.

*Radiation:*

Global (shortwave) radiation ( $G$ )	=	$200 \text{ W m}^{-2}$
Albedo	=	$0.4$
Longwave incoming radiation $L_{in}$	=	$280 \text{ W m}^{-2}$

*Other heat fluxes:*

Sensible heat flux ( $Q_H$ )	=	$30 \text{ W m}^{-2}$
Latent heat flux ( $Q_L$ )	=	$-20 \text{ W m}^{-2}$
Warming of the cold surface layer consumes		$5 \text{ W m}^{-2}$ .

Latent heat of fusion  $L_f$  =  $334,000 \text{ J kg}^{-1}$  (needed to compute **melt**)

Latent heat of vaporization  $L_{fv}$  =  $2,883,000 \text{ J kg}^{-1}$  (needed to compute **sublimation**)

**a) How much energy is available for melt  $Q_m$ ?**

$$Q_m = G - R + L_{in} - L_{out} + Q_H + Q_L + Q_{ice}$$

*(R=reflected shortwave radiation)*

Note that longwave outgoing radiation was not measured but it can be estimated from the radiation from the information given above. Note that the ice heat flux  $Q_{ice}$  is negative, if energy is consumed for warming the ice.

Net radiation ( $\text{W m}^{-2}$ ) =

Energy available for melt ( $\text{W m}^{-2}$ ) =

**b) How much melt occurs during that day in  $\text{kg m}^{-2}$  and in mm w.e. ? (Note  $1 \text{ W} = 1 \text{ J s}^{-1}$ )**

**c) How much ablation occurs during that day (in  $\text{kg m}^{-2}$  and in mm w.e.) ?**

Note that ablation includes all processes that remove mass from a glacier, i.e. also mass removed by sublimation (indicated by a negative latent heat flux) and not only melt.

## 2.) TIME SERIES

Figure 1 shows hourly time series of meteorological observations and energy balance components at a weather station on a glacier.

a) **Label what is what on the figure.** Shown are

- air temperature (K), global radiation, latent heat flux,
- longwave incoming and outgoing radiation, net radiation,
- rain heat flux, sensible heat flux,
- relative humidity (%), and wind speed ( $\text{m s}^{-1}$ ). All energy fluxes in  $\text{W m}^{-2}$ .

b) **Identify a foggy and clear-sky day** (Name at least 2 indicators).

c) **Identify a period of surface melting and a period with surface temperatures below freezing**




Figure 1. Meteorological observations and energy balance components on a glacier. Tics every 10 days.

### 3.) SPATIAL DISTRIBUTION OF ENERGY BALANCE COMPONENTS

Figure 3 shows the modeled spatial pattern of the energy fluxes on Storglaciären, Sweden, averaged over the period 7 June to 17 September 1993. The model is forced by weather station data from one station in the center of the glacier. Shown are (all energy fluxes in  $W m^{-2}$ ):

• sensible heat flux	• global radiation	• longwave incoming radiation
• latent heat flux	• diffuse radiation	• outgoing longwave radiation
• melt (in cm) over the >3 months period.	• net radiation	• direct radiation

Air temperature (not shown) is extrapolated in the model assuming a linear decrease in temperature with elevation. Humidity and wind speed (not shown) measured at the weather station were assumed constant in space. The glacier surface was at the melting point during most of the period. Cloud cover was assumed uniform across the glacier.

Which component does each subplot show ? Note that the 3 smaller maps have a different scale (you can ignore them).

### SURFACE ENERGY FLUXES ON STORGLACIÄREN Jun 7 - Sep 17, 1993




Figure 2. Calculated energy fluxes on Storglaciären, Sweden, averaged over the period 7 June to 17 September 1993. Note the smaller plots within some of the subplots show the same quantities with a different scale to better visualize any small-scale spatial variations.

**Additional exercises (optional) if time permits:**

**5.) EFFECT OF CLOUDS ON MELT**

Assume a clear sunny day on a melting glacier in summer, suddenly thick low clouds start moving in: How will net radiation change? (Use numbers for illustration, for example assume  $G=600 \text{ Wm}^{-2}$  (clear-sky),  $G=100$  or  $200 \text{ Wm}^{-2}$  (cloudy));

Longwave incoming radiation  $L_{in}$  is typically around  $250 \text{ W m}^{-2}$  during a sunny summer day, while it is around  $300 \text{ W m}^{-2}$  during foggy conditions. Longwave outgoing radiation is fixed when the surface is melting.

(Clouds affect value of albedo but this should be neglected here in your considerations; assume that the albedo is constant).

**6.) EFFECT OF RAIN ON MELT**

Assume a rainy day with  $10 \text{ mm d}^{-1}$ . Daily mean near-surface air temperature is  $10^\circ\text{C}$ . Calculate the rain heat flux and compare it to typical values of net radiation and the turbulent heat fluxes (see Figure 2 and 3).

$$Q_R = \rho_w c_w R(T_r - T_s)$$

$\rho_w$  = density of water ( $1000 \text{ kg/m}^3$ )

$c_w$  = specific heat of water,  $4180 \text{ J kg}^{-1} \text{ K}^{-1}$

$R$  = rain intensity (e.g.  $\text{mm/day}$ )

$T_r$  = rain temperature (K or Celcius)

$T_s$  = glacier surface temperature (K or Celcius); assume melting surface